

**FINAL JEE–MAIN EXAMINATION – JANUARY, 2023**  
**Held On Sunday 29th January, 2023**  
**TIME : 03:00 PM to 06:00 PM**

**SECTION-A**

61. The statement  $B \Rightarrow ((\sim A) \vee B)$  is equivalent to :
- $B \Rightarrow (A \Rightarrow B)$
  - $A \Rightarrow (A \Leftrightarrow B)$
  - $A \Rightarrow ((\sim A) \Rightarrow B)$
  - $B \Rightarrow ((\sim A) \Rightarrow B)$

**Official Ans. by NTA (1,3,4)**

**Ans. (1 or 3 or 4)**

**Sol.**

A	B	$\sim A$	$\sim A \vee B$	$B \Rightarrow ((\sim A) \vee B)$
T	T	F	T	T
T	F	F	F	T
F	T	T	T	T
F	F	T	T	T

$A \Rightarrow B$	$\sim A \Rightarrow B$	$B \Rightarrow (A \Rightarrow B)$	$A \Rightarrow ((\sim A) \Rightarrow B)$	$B \Rightarrow ((\sim A) \Rightarrow B)$
T	T	T	T	T
F	T	T	T	T
T	T	T	T	T
T	F	T	T	T

62. Shortest distance between the lines

$$\frac{x-1}{2} = \frac{y+8}{-7} = \frac{z-4}{5} \text{ and } \frac{x-1}{2} = \frac{y-2}{1} = \frac{z-6}{-3} \text{ is}$$

- $2\sqrt{3}$
- $4\sqrt{3}$
- $3\sqrt{3}$
- $5\sqrt{3}$

**Official Ans. by NTA (2)**

**Ans. (2)**

**Sol.**  $\frac{x-1}{2} = \frac{y+8}{-7} = \frac{z-4}{5} \quad \vec{a} = \hat{i} - 8\hat{j} + 4\hat{k}$

$$\frac{x-1}{2} = \frac{y-2}{1} = \frac{z-6}{-3} \quad \vec{b} = \hat{i} + 2\hat{j} + 6\hat{k}$$

$$\vec{p} = 2\hat{i} - 7\hat{j} + 5\hat{k}, \vec{q} = 2\hat{i} + \hat{j} - 3\hat{k}$$

$$\vec{p} \times \vec{q} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & -7 & 5 \\ 2 & 1 & -3 \end{vmatrix}$$

$$= \hat{i}(16) - \hat{j}(-16) + \hat{k}(16)$$

$$= 16(\hat{i} + \hat{j} + \hat{k})$$

$$d = \left| \frac{(a-b) \cdot (\vec{p} \times \vec{q})}{|\vec{p} \times \vec{q}|} \right| = \left| \frac{(-10\hat{j} - 2\hat{k}) \cdot 16(\hat{i} + \hat{j} + \hat{k})}{16\sqrt{3}} \right|$$

$$= \left| \frac{-12}{\sqrt{3}} \right| = 4\sqrt{3}$$

63. If  $\vec{a} = \hat{i} + 2\hat{k}$ ,  $\vec{b} = \hat{i} + \hat{j} + \hat{k}$ ,  $\vec{c} = 7\hat{i} - 3\hat{k} + 4\hat{k}$ ,  $\vec{r} \times \vec{b} + \vec{b} \times \vec{c} = \vec{0}$  and  $\vec{r} \cdot \vec{a} = 0$  then  $\vec{r} \cdot \vec{c}$  is equal to :

- 34
- 12
- 36
- 30

**Official Ans. by NTA (1)**

**Ans. (1)**

**Sol.**  $\vec{r} \times \vec{b} - \vec{c} \times \vec{b} = \vec{0}$

$$\Rightarrow (\vec{r} - \vec{c}) \times \vec{b} = \vec{0}$$

$$\Rightarrow \vec{r} - \vec{c} = \lambda \vec{b}$$

$$\Rightarrow \vec{r} = \vec{c} + \lambda \vec{b}$$

And given that  $\vec{r} \cdot \vec{a} = 0$

$$\Rightarrow (\vec{c} + \lambda \vec{b}) \cdot \vec{a} = 0$$

$$\Rightarrow \vec{c} \cdot \vec{a} + \lambda \vec{b} \cdot \vec{a} = 0$$

$$\Rightarrow \lambda = \frac{-\vec{c} \cdot \vec{a}}{\vec{b} \cdot \vec{a}}$$

$$\text{Now } \vec{r} \cdot \vec{c} = (\vec{c} + \lambda \vec{b}) \cdot \vec{c}$$

$$= \left( \vec{c} - \frac{\vec{c} \cdot \vec{a}}{\vec{b} \cdot \vec{a}} \vec{b} \right) \cdot \vec{c}$$

$$= |\vec{c}| - \left( \frac{\vec{c} \cdot \vec{a}}{\vec{b} \cdot \vec{a}} \right) (\vec{b} \cdot \vec{c})$$

$$= 74 - \left[ \frac{15}{3} \right] 8$$

$$= 74 - 40 = 34$$

64. Let  $S = \{w_1, w_2, \dots\}$  be the sample space associated to a random experiment. Let  $P(w_n) = \frac{P(w_{n-1})}{2}, n \geq 2$ .

Let  $A = \{2k + 3\ell; k, \ell \in \mathbb{N}\}$  and  $B = \{w_n; n \in A\}$ . Then  $P(B)$  is equal to

- |                    |                    |
|--------------------|--------------------|
| (1) $\frac{3}{32}$ | (2) $\frac{3}{64}$ |
| (3) $\frac{1}{16}$ | (4) $\frac{1}{32}$ |

**Official Ans. by NTA (2)**

**Ans. (2)**

**Sol.** Let  $P(w_1) = \lambda$  then  $P(w_2) = \frac{\lambda}{2} \dots P(w_n) = \frac{\lambda}{2^{n-1}}$

$$\text{As } \sum_{k=1}^{\infty} P(w_k) = 1 \Rightarrow \frac{\lambda}{1 - \frac{1}{2}} = 1 \Rightarrow \lambda = \frac{1}{2}$$

$$\text{So, } P(w_n) = \frac{1}{2^n}$$

$$A = \{2k + 3\ell; k, \ell \in \mathbb{N}\} = \{5, 7, 8, 9, 10, \dots\}$$

$$B = \{w_n : n \in A\}$$

$$B = \{w_5, w_7, w_8, w_9, w_{10}, w_{11}, \dots\}$$

$$A = \mathbb{N} - \{1, 2, 3, 4, 6\}$$

$$\therefore P(B) = 1 - [P(w_1) + P(w_2) + P(w_3) + P(w_4) + P(w_6)]$$

$$= 1 - \left[ \frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \frac{1}{16} + \frac{1}{64} \right]$$

$$= 1 - \frac{32+16+8+4+1}{64} = \frac{3}{64}$$

65. The value of the integral  $\int_1^2 \left( \frac{t^4+1}{t^6+1} \right) dt$  is :

- |   |
|---|
| (1) $\tan^{-1} \frac{1}{2} + \frac{1}{3} \tan^{-1} 8 - \frac{\pi}{3}$ |
| (2) $\tan^{-1} 2 - \frac{1}{3} \tan^{-1} 8 + \frac{\pi}{3}$           |
| (3) $\tan^{-1} 2 + \frac{1}{3} \tan^{-1} 8 - \frac{\pi}{3}$           |
| (4) $\tan^{-1} \frac{1}{2} - \frac{1}{3} \tan^{-1} 8 + \frac{\pi}{3}$ |

**Official Ans. by NTA (3)**

**Ans. (3)**

**Sol.**  $I = \int_1^2 \left( \frac{t^4+1}{t^6+1} \right) dt$

$$= \int_1^2 \frac{(t^4+1-t^2)+t^2}{(t^2+1)(t^4-t^2+1)} dt$$

$$= \int_1^2 \left( \frac{1}{t^2+1} + \frac{t^2}{t^6+1} \right) dt$$

$$= \int_1^2 \left( \frac{1}{t^2+1} + \frac{1}{3} \frac{3t^2}{(t^3)^2+1} \right) dt$$

$$= \tan^{-1}(t) + \frac{1}{3} \tan^{-1}(t^3) \Big|_1^2$$

$$= (\tan^{-1}(2) - \tan^{-1}(1)) + \frac{1}{3} (\tan^{-1}(2^3) - \tan^{-1}(1^3))$$

$$= \tan^{-1}(2) + \frac{1}{3} \tan^{-1}(8) - \frac{\pi}{3}$$

66. Let K be the sum of the coefficients of the odd powers of x in the expansion of  $(1+x)^{99}$ . Let a be

the middle term in the expansion of  $\left(2 + \frac{1}{\sqrt{2}}\right)^{200}$ . If

$$\frac{200}{a} C_{99} K = \frac{2^\ell m}{n}, \text{ where } m \text{ and } n \text{ are odd numbers,}$$

then the ordered pair  $(\ell, n)$  is equal to :

(1) (50, 51)

(2) (51, 99)

(3) (50, 101)

(4) (51, 101)

**Official Ans. by NTA (3)**

**Ans. (3)**

**Sol.** In the expansion of

$$(1+x)^{99} = C_0 + C_1 x + C_2 x^2 + \dots + C_{99} x^{99}$$

$$K = C_1 + C_3 + \dots + C_{99} = 2^{98}$$

a  $\Rightarrow$  Middle in the expansion of  $\left(2 + \frac{1}{\sqrt{2}}\right)^{200}$

$$T_{\frac{200}{2}+1} = {}^{200}C_{100} (2)^{100} \left(\frac{1}{\sqrt{2}}\right)^{100}$$

$$= {}^{200}C_{100} \cdot 2^{50}$$

$$\text{So, } \frac{{}^{200}C_{99} \times 2^{98}}{{}^{200}C_{100} \times 2^{50}} = \frac{100}{101} \times 2^{48}$$

$$\text{So, } \frac{25}{101} \times 2^{50} = \frac{m}{n} 2^\ell$$

$\therefore m, n$  are odd so

( $\ell, n$ ) become (50, 101) Ans.

- 67.** Let  $f$  and  $g$  be twice differentiable functions on  $\mathbb{R}$  such that

$$f''(x) = g''(x) + 6x$$

$$f'(1) = 4g'(1) - 3 = 9$$

$$f(2) = 3g(2) = 12$$

Then which of the following is NOT true ?

(1)  $g(-2) - f(-2) = 20$

(2) If  $-1 < x < 2$ , then  $|f(x) - g(x)| < 8$

(3)  $|f'(x) - g'(x)| < 6 \Rightarrow -1 < x < 1$

(4) There exists  $x_0 \in \left(1, \frac{3}{2}\right)$  such that  $f(x_0) = g(x_0)$

**Official Ans. by NTA (2)**

**Ans. (2)**

**Sol.**  $f''(x) = g''(x) + 6x \quad \dots(1)$

$$f'(1) = 4g'(1) - 3 = 9 \quad \dots(2)$$

$$f(2) = 3g(2) = 12 \quad \dots(3)$$

By integrating (1)

$$f'(x) = g'(x) + 6 \frac{x^2}{2} + C$$

At  $x = 1$ ,

$$f'(1) = g'(1) + 3 + C$$

$$\Rightarrow 9 = 4 + 3 + C \Rightarrow C = 3$$

$$\therefore f'(x) = g'(x) + 3x^2 + 3$$

Again by integrating,

$$f(x) = g(x) + \frac{3x^3}{3} + 3x + D$$

At  $x = 2$ ,

$$f(2) = g(2) + 8 + 3(2) + D$$

$$\Rightarrow 12 = 4 + 8 + 6 + D \Rightarrow D = -6$$

$$\text{So, } f(x) = g(x) + x^3 + 3x - 6$$

$$\Rightarrow f(x) - g(x) = x^3 + 3x - 6$$

At  $x = -2$ ,

$$\Rightarrow g(-2) - f(-2) = 20 \quad (\text{Option (1) is true})$$

Now, for  $-1 < x < 2$

$$h(x) = f(x) - g(x) = x^3 + 3x - 6$$

$$\Rightarrow h'(x) = 3x^2 + 3$$

$$\Rightarrow h(x) \uparrow$$

$$\text{So, } h(-1) < h(x) < h(2)$$

$$\Rightarrow -10 < h(x) < 8$$

$$\Rightarrow |h(x)| < 10 \quad (\text{option (2) is NOT true})$$

$$\text{Now, } h'(x) = f'(x) - g'(x) = 3x^2 + 3$$

$$\text{If } |h'(x)| < 6 \Rightarrow |3x^2 + 3| < 6$$

$$\Rightarrow 3x^2 + 3 < 6$$

$$\Rightarrow x^2 < 1$$

$$\Rightarrow -1 < x < 1 \quad (\text{option (3) is True})$$

$$\text{If } x \in (-1, 1) \quad |f'(x) - g'(x)| < 6$$

option (3) is true and now to solve

$$f(x) - g(x) = 0$$

$$\Rightarrow x^3 + 3x - 6 = 0$$

$$h(x) = x^3 + 3x - 6$$

$$\text{here, } h(1) = -\text{ve} \text{ and } h\left(\frac{3}{2}\right) = +\text{ve}$$

$$\text{So there exists } x_0 \in \left(1, \frac{3}{2}\right) \text{ such that } f(x_0) = g(x_0)$$

(option (4) is true)

68. The set of all values of  $t \in \mathbb{R}$ , for which the matrix

$$\begin{bmatrix} e^t & e^{-t}(\sin t - 2\cos t) & e^{-t}(-2\sin t - \cos t) \\ e^t & e^{-t}(2\sin t + \cos t) & e^{-t}(\sin t - 2\cos t) \\ e^t & e^{-t}\cos t & e^{-t}\sin t \end{bmatrix} \quad \text{is}$$

invertible, is

(1)  $\left\{(2k+1)\frac{\pi}{2}, k \in \mathbb{Z}\right\}$     (2)  $\left\{k\pi + \frac{\pi}{4}, k \in \mathbb{Z}\right\}$

(3)  $\{k\pi, k \in \mathbb{Z}\}$     (4)  $\mathbb{R}$

**Official Ans. by NTA (4)**

**Ans. (4)**

- Sol. If its invertible, then determinant value  $\neq 0$

So,

$$\begin{vmatrix} e^t & e^{-t}(\sin t - 2\cos t) & e^{-t}(-2\sin t - \cos t) \\ e^t & e^{-t}(2\sin t + \cos t) & e^{-t}(\sin t - 2\cos t) \\ e^t & e^{-t}\cos t & e^{-t}\sin t \end{vmatrix} \neq 0$$

$$\Rightarrow e^t \cdot e^{-t} \cdot e^{-t} \begin{vmatrix} 1 & \sin t - 2\cos t & -2\sin t - \cos t \\ 1 & 2\sin t + \cos t & \sin t - 2\cos t \\ 1 & \cos t & \sin t \end{vmatrix} \neq 0$$

Applying,  $R_1 \rightarrow R_1 - R_2$  then  $R_2 \rightarrow R_2 - R_3$

We get

$$e^{-t} \begin{vmatrix} 0 & -\sin t - \cos t & -3\sin t + \cos t \\ 0 & 2\sin t & -2\cos t \\ 1 & \cos t & \sin t \end{vmatrix} \neq 0$$

By expanding we have,

$$e^{-t} \times 1 \left( 2\sin t \cos t + 6\cos^2 t + 6\sin^2 t - 2\sin t \cos t \right) \neq 0$$

$$\Rightarrow e^{-t} \times 6 \neq 0$$

for  $\forall t \in \mathbb{R}$

69. The area of the region

$$A = \left\{ (x, y) : |\cos x - \sin x| \leq y \leq \sin x, 0 \leq x \leq \frac{\pi}{2} \right\}$$

(1)  $1 - \frac{3}{\sqrt{2}} + \frac{4}{\sqrt{5}}$     (2)  $\sqrt{5} + 2\sqrt{2} - 4.5$

(3)  $\frac{3}{\sqrt{5}} - \frac{3}{\sqrt{2}} + 1$     (4)  $\sqrt{5} - 2\sqrt{2} + 1$

**Official Ans. by NTA (4)**

**Ans. (4)**


- Sol.**  $|\cos x - \sin x| \leq y \leq \sin x$

Intersection point of  $\cos x - \sin x = \sin x$

$$\Rightarrow \tan x = \frac{1}{2}$$

$$\text{Let } \psi = \tan^{-1} \frac{1}{2}$$

$$\text{So, } \tan \psi = \frac{1}{2}, \sin \psi = \frac{1}{\sqrt{5}}, \cos \psi = \frac{2}{\sqrt{5}}$$



$$\text{Area} = \int_{\psi}^{\pi/2} (\sin x - |\cos x - \sin x|) dx$$

$$= \int_{\psi}^{\pi/4} (\sin x - (\cos x - \sin x)) dx$$

$$+ \int_{\pi/4}^{\pi/2} (\sin x - (\sin x - \cos x)) dx$$

$$= \int_{\psi}^{\pi/4} (2\sin x - \cos x) dx + \int_{\pi/4}^{\pi/2} \cos x dx$$

$$= [-2\cos x - \sin x]_{\psi}^{\pi/4} + [\sin x]_{\pi/4}^{\pi/2}$$

$$= -\sqrt{2} - \frac{1}{\sqrt{2}} + 2\cos \psi + \sin \psi + \left(1 - \frac{1}{\sqrt{2}}\right)$$

$$= -\sqrt{2} - \frac{1}{\sqrt{2}} + 2\left(\frac{2}{\sqrt{5}}\right) + \left(\frac{1}{\sqrt{5}}\right) + 1 - \frac{1}{\sqrt{2}}$$

$$= \sqrt{5} - 2\sqrt{2} + 1$$

70. The set of all values of  $\lambda$  for which the equation  $\cos^2 2x - 2\sin^4 x - 2\cos^2 x = \lambda$

(1)  $[-2, -1]$     (2)  $\left[-2, -\frac{3}{2}\right]$

(3)  $\left[-1, -\frac{1}{2}\right]$     (4)  $\left[-\frac{3}{2}, -1\right]$

**Official Ans. by NTA (4)**

**Ans. (4)**







79. Let  $R$  be a relation defined on  $\mathbb{N}$  as  $a R b$  if  $2a + 3b$  is a multiple of 5,  $a, b \in \mathbb{N}$ . Then  $R$  is
- not reflexive
  - transitive but not symmetric
  - symmetric but not transitive
  - an equivalence relation

**Official Ans. by NTA (4)**
**Ans. (4)**
**Sol.**  $a R a \Rightarrow 5a$  is multiple of 5

So reflexive

$$a R b \Rightarrow 2a + 3b = 5\alpha,$$

 Now  $b R a$ 

$$\begin{aligned} 2b + 3a &= 2b + \left(\frac{5\alpha - 3b}{2}\right) \cdot 3 \\ &= \frac{15}{2}\alpha - \frac{5}{2}b = \frac{5}{2}(3\alpha - b) \\ &= \frac{5}{2}(2a + 2b - 2\alpha) \\ &= 5(a + b - \alpha) \end{aligned}$$

Hence symmetric

$$a R b \Rightarrow 2a + 3b = 5\alpha.$$

$$b R c \Rightarrow 2b + 3c = 5\beta$$

$$\text{Now } 2a + 5b + 3c = 5(\alpha + \beta)$$

$$\Rightarrow 2a + 5b + 3c = 5(\alpha + \beta)$$

$$\Rightarrow 2a + 3c = 5(\alpha + \beta - b)$$

$$\Rightarrow a R c$$

Hence relation is equivalence relation.

80. Consider a function  $f: \mathbb{N} \rightarrow \mathbb{R}$ , satisfying

$$f(1) + 2f(2) + 3f(3) + \dots + xf(x) = x(x+1)f(x); x \geq 2$$

with  $f(1)=1$ . Then  $\frac{1}{f(2022)} + \frac{1}{f(2028)}$  is equal to

- 8200
- 8000
- 8400
- 8100

**Official Ans. by NTA (4)**
**Ans. (4)**
**Sol.** Given for  $x \geq 2$ 

$$f(1) + 2f(2) + \dots + xf(x) = x(x+1)f(x)$$

 replace  $x$  by  $x+1$ 

$$\Rightarrow x(x+1)f(x) + (x+1)f(x+1)$$

$$= (x+1)(x+2)f(x+1)$$

$$\Rightarrow \frac{x}{f(x+1)} + \frac{1}{f(x)} = \frac{(x+2)}{f(x)}$$

$$\Rightarrow x f(x) = (x+1) f(x+1) = \frac{1}{2}, x \geq 2$$

$$f(2) = \frac{1}{4}, f(3) = \frac{1}{6}$$

$$\text{Now } f(2022) = \frac{1}{4044}$$

$$f(2028) = \frac{1}{4056}$$

$$\text{So, } \frac{1}{f(2022)} + \frac{1}{f(2028)} = 4044 + 4056 = 8100$$

## SECTION-B

81. The total number of 4-digit numbers whose greatest common divisor with 54 is 2, is \_\_\_\_\_.

**Official Ans. by NTA (3000)**
**Ans. (3000)**
**Sol.**  $N$  should be divisible by 2 but not by 3

$$N = (\text{Numbers divisible by 2}) - (\text{Numbers divisible by 6})$$

$$N = \frac{9000}{2} - \frac{9000}{6} = 4500 - 1500 = 3000$$

82. A triangle is formed by the tangents at the point  $(2, 2)$  on the curves  $y^2 = 2x$  and  $x^2 + y^2 = 4x$ , and the line  $x + y + 2 = 0$ . If  $r$  is the radius of its circumcircle, then  $r^2$  is equal to \_\_\_\_\_.

**Official Ans. by NTA (10)**
**Ans. (10)**

**Sol.**  $S_1 : y^2 = 2x$        $S_2 : x^2 + y^2 = 4x$

$P(2,2)$  is common point on  $S_1$  &  $S_2$


$$T_1 \text{ is tangent to } S_1 \text{ at } P \Rightarrow T_1 : y \cdot 2 = x + 2$$

$$\Rightarrow T_1 : x - 2y + 2 = 0$$

$$T_2 \text{ is tangent to } S_2 \text{ at } P \Rightarrow T_2 : x \cdot 2 + y \cdot 2 = 2(x+2)$$

$$\Rightarrow T_2 : y = 2$$

&  $L_3 : x + y + 2 = 0$  is third line



$$PQ = a = \sqrt{20}$$

$$QR = b = \sqrt{8}$$

$$RP = c = 6$$

$$\text{Area } (\Delta PQR) = \Delta = \frac{1}{2} \times 6 \times 2 = 6$$

$$\therefore r = \frac{abc}{4\Delta} = \frac{\sqrt{160}}{4} = \sqrt{10} \Rightarrow r^2 = 10$$

- 83.** A circle with centre  $(2, 3)$  and radius 4 intersects the line  $x + y = 3$  at the points P and Q. If the tangents at P and Q intersect at the point  $S(\alpha, \beta)$ , then  $4\alpha - 7\beta$  is equal to \_\_\_\_\_.

**Official Ans. by NTA (11)**

**Ans. (11)**

**Sol.** The given line is polar or  $P(2, \beta)$  w.r.t. given circle

$$x^2 + y^2 - 4x - 6y - 3 = 0$$

Chord or contact

$$\alpha x + \beta y - 2(x + \alpha) - 3(y + \beta) - 3 = 0$$

$$\Rightarrow (\alpha - 2)x + (\beta - 3)y - (2\alpha + 3\beta + 3) = 0 \dots\dots (i)$$

$\because$  But the equation of chord of contact is given

$$\text{as : } x + y - 3 = 0 \dots\dots (ii)$$

comparing the coefficients

$$\frac{\alpha - 2}{1} = \frac{\beta - 3}{1} = -\left(\frac{2\alpha + 3\beta + 3}{-3}\right)$$

$$\text{On solving } \alpha = -6$$

$$\beta = -5$$

$$\text{Now } 4\alpha - 7\beta = 11$$

- 84.** Let  $a_1 = b_1 = 1$  and  $a_n = a_{n-1} + (n - 1)$ ,  $b_n = b_{n-1} + a_{n-1}$ ,  $\forall n \geq 2$ . If  $S = \sum_{n=1}^{10} \frac{b_n}{2^n}$  and  $T = \sum_{n=1}^8 \frac{n}{2^{n-1}}$ , then  $2^7(2S - T)$  is equal to \_\_\_\_\_.

**Official Ans. by NTA (461)**

**Ans. (461)**

**Sol.** As,  $S = \frac{b_1}{2} + \frac{b_2}{2^2} + \dots\dots + \frac{b_9}{2^9} + \frac{b_{10}}{2^{10}}$

$$\Rightarrow \frac{S}{2} = \frac{b_1}{2^2} + \frac{b_2}{2^3} + \dots\dots + \frac{b_9}{2^{10}} + \frac{b_{10}}{2^{11}}$$

subtracting

$$\Rightarrow \frac{S}{2} = \frac{b_1}{2} + \left( \frac{a_1}{2^2} + \frac{a_2}{2^3} + \dots\dots + \frac{a_9}{2^{10}} \right) - \frac{b_{10}}{2^{11}}$$

$$\Rightarrow S = b_1 - \frac{b_{10}}{2^{10}} + \left( \frac{a_1}{2} + \frac{a_2}{2^2} + \dots\dots + \frac{a_9}{2^9} \right)$$

$$\Rightarrow \frac{S}{2} = \frac{b_1}{2} - \frac{b_{10}}{2^{11}} + \left( \frac{a_1}{2^2} + \frac{a_2}{2^3} + \dots\dots + \frac{a_9}{2^{10}} \right)$$

subtracting

$$\Rightarrow \frac{S}{2} = \frac{b_1}{2} - \frac{b_{10}}{2^{11}} + \left( \frac{a_1}{2} - \frac{a_9}{2^{10}} \right) + \left( \frac{1}{2^2} + \frac{2}{2^3} + \dots + \frac{8}{2^9} \right)$$

$$\Rightarrow \frac{S}{2} = \frac{a_1 + b_1}{2} - \frac{(b_{10} + 2a_9)}{2^{11}} + \frac{T}{4}$$

$$\Rightarrow 2S = 2(a_1 + b_1) - \frac{(b_{10} + 2a_9)}{2^9} + T$$

$$\Rightarrow 2^7 (2S - T) = 2^8 (a_1 + b_1) - \frac{(b_{10} + 2a_9)}{4}$$

Given  $a_n - a_{n-1} = n - 1$ ,

$$\therefore a_2 - a_1 = 1$$

$$a_3 - a_2 = 2$$

⋮

$$a_9 - a_8 = 8$$

$$a_9 - a_1 = 1 + 2 + \dots + 8 = 36$$

$$\Rightarrow a_9 = 37 \quad (a_1 = 1)$$

Also,  $b_n - b_{n-1} = a_{n-1}$

$$\therefore b_{10} - b_1 = a_1 + a_2 + \dots + a_9$$

$$= 1 + 2 + 4 + 7 + 11 + 16 + 22 + 29 + 37$$

$$\Rightarrow b_{10} = 130 \quad (\text{As } b_1 = 1)$$

$$\therefore 2^7 (2S - T) = 2^8 (1 + 1) - (130 + 2 \times 37)$$

$$2^9 - \frac{204}{4} = 461$$

- 85.** If the equation of the normal to the curve

$$y = \frac{x-a}{(x+b)(x-2)}$$
 at the point  $(1, -3)$  is  $x - 4y = 13$ ,

then the value of  $a + b$  is equal to \_\_\_\_.

**Official Ans. by NTA (4)**

**Ans. (4)**

**Sol.**  $y = \frac{x-a}{(x+b)(x-2)}$

At point  $(1, -3)$ ,

$$-3 = \frac{1-9}{(1+b)(1-2)}$$

$$\Rightarrow 1 - a = 3(1 + b) \quad \dots \dots (1)$$

Now,  $y = \frac{x-a}{(x+b)(x-2)}$

$$\Rightarrow \frac{dy}{dx} = \frac{(x+b)(x-2) \times (1) - (x-a)(2x+b-2)}{(x+b)^2(x-2)^2}$$

At  $(1, -3)$  slope of normal is  $\frac{1}{4}$  hence  $\frac{dy}{dx} = -4$ ,

$$\text{So, } -4 = \frac{(1+b)(-1) - (1-a)b}{(1+b)^2(-1)^2}$$

Using equation (1)

$$\Rightarrow -4 = \frac{(1+b)(-1) - 3(b+1)b}{(1+b)^2}$$

$$\Rightarrow -4 = \frac{(-1) - 3b}{(1+b)} \quad (b \neq -1)$$

$$\Rightarrow b = -3$$

$$\text{So, } a = 7$$

$$\text{Hence, } a + b = 7 - 3 = 4$$

- 86.** Let  $A$  be a symmetric matrix such that  $|A| = 2$  and

$$\begin{bmatrix} 2 & 1 \\ 3 & \frac{3}{2} \end{bmatrix} A = \begin{bmatrix} 1 & 2 \\ \alpha & \beta \end{bmatrix}$$
. If the sum of the diagonal elements of  $A$  is  $s$ , then  $\frac{\beta s}{\alpha^2}$  is equal to \_\_\_\_.

**Official Ans. by NTA (5)**

**Ans. (5)**

**Sol.**  $\begin{bmatrix} 2 & 1 \\ 3 & \frac{3}{2} \end{bmatrix} \begin{bmatrix} a & b \\ b & c \end{bmatrix} = \begin{bmatrix} 1 & 2 \\ \alpha & \beta \end{bmatrix}$

$$\text{Now } ac - b^2 = 2 \quad \text{and} \quad 2a + b = 1$$

$$\text{and } 2b + c = 2$$

solving all these above equations we get

$$\frac{1-b}{2} \times \left( \frac{2-2b}{1} \right) - b^2 = 2$$

$$\Rightarrow (1-b)^2 - b^2 = 2$$

$$\Rightarrow 1 - 2b = 2$$

$$\Rightarrow b = -\frac{1}{2} \quad \text{and} \quad a = \frac{3}{4} \quad \text{and} \quad c = 3$$

$$\text{Hence } \alpha = 3a + \frac{3b}{2} = \frac{9}{4} - \frac{3}{4} = \frac{3}{2}$$

$$\text{and } \beta = 3b + \frac{3c}{2} = -\frac{3}{2} + \frac{9}{2} = 3$$

$$\text{also } s = a + c = \frac{15}{4}$$

$$\therefore \frac{\beta s}{\alpha^2} = \frac{3 \times 15}{4 \times \frac{9}{4}} = 5$$

87. Let  $\{a_k\}$  and  $\{b_k\}$ ,  $k \in \mathbb{N}$ , be two G.P.s with common ratio  $r_1$  and  $r_2$  respectively such that  $a_1 = b_1 = 4$  and  $r_1 < r_2$ . Let  $c_k = a_k + b_k$ ,  $k \in \mathbb{N}$ .

If  $c_2 = 5$  and  $c_3 = \frac{13}{4}$  then  $\sum_{k=1}^{\infty} c_k - (12a_6 + 8b_4)$  is equal to \_\_\_\_\_.

**Official Ans. by NTA (9)**

**Ans. (9)**

**Sol.** Given that

$$\begin{aligned} c_k &= a_k + b_k \text{ and } a_1 = b_1 = 4 \\ \text{also } a_2 &= 4r_1 & a_3 &= 4r_1^2 \\ b_2 &= 4r_2 & b_3 &= 4r_2^2 \end{aligned}$$

$$\text{Now } c_2 = a_2 + b_2 = 5 \text{ and } c_3 = a_3 + b_3 = \frac{13}{4}$$

$$\Rightarrow r_1 + r_2 = \frac{5}{4} \text{ and } r_1^2 + r_2^2 = \frac{13}{16}$$

$$\text{Hence } r_1 r_2 = \frac{3}{8} \text{ which gives } r_1 = \frac{1}{2} \text{ & } r_2 = \frac{3}{4}$$

$$\begin{aligned} \sum_{k=1}^{\infty} c_k - (12a_6 + 8b_4) &= \frac{4}{1-r_1} + \frac{4}{1-r_2} - \left( \frac{48}{32} + \frac{27}{2} \right) \\ &= 24 - 15 = 9 \end{aligned}$$

88. Let  $X = \{11, 12, 13, \dots, 40, 41\}$  and  $Y = \{61, 62, 63, \dots, 90, 91\}$  be the two sets of observations. If  $\bar{x}$  and  $\bar{y}$  are their respective means and  $\sigma^2$  is the variance of all the observations in  $X \cup Y$ , then  $|\bar{x} + \bar{y} - \sigma^2|$  is equal to \_\_\_\_\_.

**Official Ans. by NTA (603)**

**Ans. (603)**

**Sol.**  $\bar{x} = \frac{\sum_{i=11}^{41} i}{31} = \frac{11+41}{2} = 26$  (31 elements)

$$\bar{y} = \frac{\sum_{j=61}^{91} j}{31} = \frac{61+91}{2} = 76$$
 (31 elements)

$$\text{Combined mean, } \mu = \frac{31 \times 26 + 31 \times 76}{31+31}$$

$$= \frac{26+76}{2} = 51$$

$$\sigma^2 = \frac{1}{62} \times \left( \sum_{i=1}^{31} (x_i - \mu)^2 + \sum_{i=1}^{31} (y_i - \mu)^2 \right) = 705$$

Since,  $x_i \in X$  are in A.P. with 31 elements & common difference 1, same is  $y_i \in Y$ , when written in increasing order.

$$\begin{aligned} \therefore \sum_{i=1}^{31} (x_i - \mu)^2 &= \sum_{i=1}^{31} (y_i - \mu)^2 \\ &= 10^2 + 11^2 + \dots + 40^2 \\ &= \frac{40 \times 41 \times 81}{6} - \frac{9 \times 10 \times 19}{6} = 21855 \\ \therefore |\bar{x} + \bar{y} - \sigma^2| &= |26 + 76 - 705| = 603 \end{aligned}$$

89. Let  $\alpha = 8 - 14i$ ,  $A = \left\{ z \in \mathbb{C} : \frac{\alpha z - \bar{\alpha} \bar{z}}{z^2 - (\bar{z})^2 - 112i} = 1 \right\}$

and  $B = \left\{ z \in \mathbb{C} : |z + 3i| = 4 \right\}$ .

Then  $\sum_{z \in A \cap B} (Re z - Im z)$  is equal to \_\_\_\_\_.

**Official Ans. by NTA (14)**

**Ans. (14)**

**Sol.**  $\alpha = 8 - 14i$

$$z = x + iy$$

$$az = (8x + 14y) + i(-14x + 8y)$$

$$z + \bar{z} = 2x \quad z - \bar{z} = 2iy$$

$$\text{Set A: } \frac{2i(-14x + 8y)}{i(4xy - 112)} = 1$$

$$(x - 4)(y + 7) = 0$$

$$x = 4 \quad \text{or} \quad y = -7$$

$$\text{Set B: } x^2 + (y + 3)^2 = 16$$

$$\text{when } x = 4 \quad y = -3$$

$$\text{when } y = -7 \quad x = 0$$

$$\therefore A \cap B = \{4 - 3i, 0 - 7i\}$$

$$\text{So, } \sum_{z \in A \cap B} (\operatorname{Re} z - \operatorname{Im} z) = 4 - (-3) + (0 - (-7)) = 14$$

- 90.** Let  $\alpha_1, \alpha_2, \dots, \alpha_7$  be the roots of the equation  $x^7 + 3x^5 - 13x^3 - 15x = 0$  and  $|\alpha_1| \geq |\alpha_2| \geq \dots \geq |\alpha_7|$ .

Then  $\alpha_1\alpha_2 - \alpha_3\alpha_4 + \alpha_5\alpha_6$  is equal to \_\_\_\_\_.

**Official Ans. by NTA (9)**

**Ans. (9)**

**Sol.** Given equation can be rearranged as

$$x(x^6 + 3x^4 - 13x^2 - 15) = 0$$

clearly  $x = 0$  is one of the root and other part can be observed by replacing  $x^2 = t$  from which we have

$$t^3 + 3t^2 - 13t - 15 = 0$$

$$\Rightarrow (t - 3)(t^2 + 6t + 5) = 0$$

$$\text{So, } t = 3, t = -1, t = -5$$

Now we are getting  $x^2 = 3, x^2 = -1, x^2 = -5$

$$\Rightarrow x = \pm\sqrt{3}, x = \pm i, x = \pm\sqrt{5}i$$

From the given condition  $|\alpha_1| \geq |\alpha_2| \geq \dots \geq |\alpha_7|$

We can clearly say that  $|\alpha_7| = 0$  and

$$\text{and } |\alpha_6| = \sqrt{5} = |\alpha_5|$$

$$\text{and } |\alpha_4| = \sqrt{3} = |\alpha_3| \text{ and } |\alpha_2| = 1 = |\alpha_1|$$

So we can have,  $\alpha_1 = \sqrt{5}i, \alpha_2 = -\sqrt{5}i, \alpha_3 = \sqrt{3}i,$

$$\alpha_4 = -\sqrt{3}, \alpha_5 = i, \alpha_6 = -i$$

Hence

$$\alpha_1\alpha_2 - \alpha_3\alpha_4 + \alpha_5\alpha_6$$

$$= 1 - (-3) + 5 = 9$$