

**FINAL JEE–MAIN EXAMINATION – JANUARY, 2023**  
**Held On Monday 30th January, 2023**  
**TIME : 09:00 AM to 12:00 PM**

**SECTION-A**

- 61.** Let  $A = \begin{bmatrix} m & n \\ p & q \end{bmatrix}$ ,  $d = |A| \neq 0$   $|A - d(\text{adj } A)| = 0$ . Then

(1)  $(1+d)^2 = (m+q)^2$

(2)  $1+d^2 = (m+q)^2$

(3)  $(1+d)^2 = m^2 + q^2$

(4)  $1+d^2 = m^2 + q^2$

**Official Ans. by NTA (1)**

**Ans. (1)**

**Sol.**  $A = \begin{bmatrix} m & n \\ p & q \end{bmatrix}$ ,  $|A - d(\text{adj } A)| = 0$

$$\Rightarrow |A - d(\text{adj } A)| = \left| \begin{bmatrix} m & n \\ p & q \end{bmatrix} - d \begin{bmatrix} q & -n \\ -p & m \end{bmatrix} \right| = \left| \begin{matrix} m - qd & n(1+d) \\ p(1+d) & q - md \end{matrix} \right| = 0$$

$$\Rightarrow (m - qd)(q - md) - np(1 + d)^2 = 0$$

$$\Rightarrow mq - m^2d - q^2d + mqd^2 - np(1 + d)^2 = 0$$

$$\Rightarrow (mq - np) + d^2(mq - np) - d(m^2 + q^2 + 2np) = 0$$

$$\Rightarrow d + d^3 - d((m+q)^2 - 2d) = 0$$

$$\Rightarrow 1 + d^2 = (m+q)^2 - 2d$$

$$\Rightarrow (1+d)^2 = (m+q)^2$$

$\therefore$  Option (1) is correct.

- 62.** The line  $l_1$  passes through the point  $(2,6,2)$  and is perpendicular to the plane  $2x + y - 2z = 10$ . Then the shortest distance between the line  $l_1$  and the line

$$\frac{x+1}{2} = \frac{y+4}{-3} = \frac{z}{2} \text{ is :}$$

(1) 7

(2)  $\frac{19}{3}$

(3)  $\frac{19}{3}$

(4) 9

**Official Ans. by NTA (4)**


**Ans. (4)**

**Sol.** Line  $\ell$ , is given by

$$L_1: \frac{x-2}{2} = \frac{y-6}{1} = \frac{z-2}{-2}$$

Given,

$$L_2: \frac{x+1}{2} = \frac{y+4}{-3} = \frac{z}{2}$$



$$\text{Shortest distance} = \frac{|\vec{AB} \cdot \vec{MN}|}{\vec{MN}}$$

$$\vec{AB} = 3\hat{i} + 10\hat{j} + 2\hat{k}$$

$$\vec{MN} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & 1 & -2 \\ 2 & -3 & 2 \end{vmatrix} = -4\hat{i} - 8\hat{j} - 8\hat{k}$$

$$MN = \sqrt{16 + 64 + 64} = 12$$

$$\therefore \text{Shortest distance} = \frac{|-12 - 80 - 16|}{12} = 9$$

$\therefore$  Option (4) is correct.

- 63.** If an unbiased die, marked with  $-2, -1, 0, 1, 2, 3$  on its faces, is thrown five times, then the probability that the product of the outcomes is positive, is :

(1)  $\frac{881}{2592}$

(2)  $\frac{521}{2592}$

(3)  $\frac{440}{2592}$

(4)  $\frac{27}{288}$

**Official Ans. by NTA (2)**

**Ans. (2)**

**Sol.** Either all outcomes are positive or any two are negative.

$$\text{Now, } p = P(\text{positive}) = \frac{3}{6} = \frac{1}{2}$$

$$q = p(\text{negative}) = \frac{2}{6} = \frac{1}{3}$$

Required probability

$$\begin{aligned} &= {}^5C_5 \left(\frac{1}{2}\right)^5 + {}^5C_2 \left(\frac{1}{3}\right)^2 \left(\frac{1}{2}\right)^3 + {}^5C_4 \left(\frac{1}{3}\right)^4 \left(\frac{1}{2}\right)^1 \\ &= \frac{521}{2592} \end{aligned}$$

$\therefore$  Option (2) is correct.

**64.** Let the system of linear equations

$$x + y + kz = 2$$

$$2x + 3y - z = 1$$

$$3x + 4y + 2z = k$$

have infinitely many solutions. Then the system

$$(k+1)x + (2k-1)y = 7$$

$$(2k+1)x + (k+5)y = 10 \text{ has :}$$

(1) infinitely many solutions

(2) unique solution satisfying  $x - y = 1$

(3) no solution

(4) unique solution satisfying  $x + y = 1$

**Official Ans. by NTA (4)**

**Ans. (4)**

$$\begin{vmatrix} 1 & 1 & k \\ 2 & 3 & -1 \\ 3 & 4 & 2 \end{vmatrix} = 0$$

$$\Rightarrow 1(10) - 1(7) + k(-1) = 0$$

$$\Rightarrow k = 3$$

For  $k = 3$ , 2<sup>nd</sup> system is

$$4x + 5y = 7 \quad \dots\dots(1)$$

$$\text{and } 7x + 8y = 10 \quad \dots\dots(2)$$

Clearly, they have a unique solution

$$(2) - (1) \Rightarrow 3x + 3y = 3$$

$$\Rightarrow x + y = 1$$

**65.** If  $\tan 15^\circ + \frac{1}{\tan 75^\circ} + \frac{1}{\tan 105^\circ} + \tan 195^\circ = 2a$ , then the value of  $\left(a + \frac{1}{a}\right)$  is :

$$(1) 4 \quad (2) 4 - 2\sqrt{3}$$

$$(3) 2 \quad (4) 5 - \frac{3}{2}\sqrt{3}$$

**Official Ans. by NTA (1)**

**Ans. (1)**

**Sol.** Option (1)

$$\tan 15^\circ = 2 - \sqrt{3}$$

$$\frac{1}{\tan 75^\circ} = \cot 75^\circ = 2 - \sqrt{3}$$

$$\frac{1}{\tan 105^\circ} = \cot(105^\circ) = -\cot 75^\circ = \sqrt{3} - 2$$

$$\tan 195^\circ = \tan 15^\circ = 2 - \sqrt{3}$$

$$\therefore 2(2 - \sqrt{3}) = 2a \Rightarrow a = 2 - \sqrt{3}$$

$$\Rightarrow a + \frac{1}{a} = 4$$

**66.** Suppose  $f : R \rightarrow (0, \infty)$  be a differentiable function such that  $5f(x+y) = f(x) \cdot f(y), \forall x, y \in R$ . If

$$f(3) = 320, \text{ then } \sum_{n=0}^5 f(n) \text{ is equal to :}$$

$$(1) 6875 \quad (2) 6575$$

$$(3) 6825 \quad (4) 6528$$

**Official Ans. by NTA (3)**

**Ans. (3)**

**Sol.** Option (3)

$$5f(x+y) = f(x) \cdot f(y)$$

$$5f(0) = f(0)^2 \Rightarrow f(0) = 5$$

$$5f(x+1) = f(x) \cdot f(1)$$

$$\Rightarrow \frac{f(x+1)}{f(x)} = \frac{f(1)}{5}$$

$$\Rightarrow \frac{f(1)}{f(0)} \cdot \frac{f(2)}{f(1)} \cdot \frac{f(3)}{f(2)} = \left(\frac{f(1)}{5}\right)^3$$

$$\Rightarrow \frac{320}{5} = \frac{(f(1))^3}{5^3} \Rightarrow f(1) = 20$$

$$\therefore 5f(x+1) = 20 \cdot f(x) \Rightarrow f(x+1) = 4f(x)$$

$$\sum_{n=0}^5 f(n) = 5 + 5 \cdot 4 + 5 \cdot 4^2 + 5 \cdot 4^3 + 5 \cdot 4^4 + 5 \cdot 4^5$$

$$= \frac{5[4^6 - 1]}{3} = 6825$$

- 67.** If  $a_n = \frac{-2}{4n^2 - 16n + 15}$ , then  $a_1 + a_2 + \dots + a_{25}$  is equal to :

- (1)  $\frac{51}{144}$       (2)  $\frac{49}{138}$   
(3)  $\frac{50}{141}$       (4)  $\frac{52}{147}$

**Official Ans. by NTA (3)**

**Ans. (3)**

**Sol.** Option (3)

If  $a_n = \frac{-2}{4n^2 - 16n + 15}$  then  $a_1 + a_2 + \dots + a_{25}$

$$\begin{aligned} \Rightarrow \sum_{n=1}^{25} a_n &= \sum \frac{-2}{4n^2 - 16n + 15} \\ &= \sum \frac{-2}{4n^2 - 6n - 10n + 15} \\ &= \sum \frac{-2}{2n(2n-3) - 5(2n-3)} \\ &= \sum \frac{-2}{(2n-3)(2n-5)} \\ &= \sum \frac{1}{2n-3} - \frac{1}{2n-5} \\ &= \frac{1}{47} - \frac{1}{(-3)} \\ &= \frac{50}{141} \end{aligned}$$

- 68.** If the coefficient of  $x^{15}$  in the expansion of  $\left(ax^3 + \frac{1}{bx^3}\right)^{15}$  is equal to the coefficient of  $x^{-15}$  in the expansion of  $\left(ax^{\frac{1}{3}} - \frac{1}{bx^3}\right)^{15}$ , where  $a$  and  $b$  are positive real numbers, then for each such ordered pair  $(a, b)$  :
- (1)  $a = b$       (2)  $ab = 1$   
(3)  $a = 3b$       (4)  $ab = 3$

**Official Ans. by NTA (2)**

**Ans. (2)**

**Sol.** Option (2)

Coefficient Of  $x^{15}$  in  $\left(ax^3 + \frac{1}{bx^3}\right)^{15}$

$$T_{r+1} = {}^{15}C_r \left(ax^3\right)^{15-r} \left(\frac{1}{bx^3}\right)^r$$

$$45 - 3r - \frac{r}{3} = 15$$

$$30 = \frac{10r}{3}$$

$$r = 9$$

Coefficient of  $x^{15} = {}^{15}C_9 a^6 b^{-9}$

Coefficient of  $x^{-15}$  in  $\left(ax^{\frac{1}{3}} - \frac{1}{bx^3}\right)^{15}$

$$T_{r+1} = {}^{15}C_r \left(ax^{\frac{1}{3}}\right)^{15-r} \left(-\frac{1}{bx^3}\right)^r$$

$$5 - \frac{r}{3} - 3r = -15$$

$$\frac{10r}{3} = 20$$

$$r = 6$$

Coefficient =  ${}^{15}C_6 a^9 \times b^{-6}$

$$\Rightarrow \frac{a^9}{b^6} = \frac{a^6}{b^9}$$

$$\Rightarrow a^3 b^3 = 1 \Rightarrow ab = 1$$

- 69.** If  $\vec{a}, \vec{b}, \vec{c}$  are three non-zero vectors and  $\hat{n}$  is a unit vector perpendicular to  $\vec{c}$  such that  $\vec{a} = \alpha \vec{b} - \hat{n}$ , ( $\alpha \neq 0$ ) and  $\vec{b} \cdot \vec{c} = 12$ , then

$|\vec{c} \times (\vec{a} \times \vec{b})|$  is equal to :

- (1) 15  
(2) 9  
(3) 12  
(4) 6

**Official Ans. by NTA (3)**

**Ans. (3)**

$$\mathbf{n} \perp \vec{c} \quad \vec{a} = \alpha \vec{b} - \vec{n}$$

$$\vec{b} \cdot \vec{c} = 12$$

$$\vec{a} \cdot \vec{c} = \alpha(\vec{b} \cdot \vec{c}) - \vec{n} \cdot \vec{c}$$

$$\vec{a} \cdot \vec{c} = \alpha(\vec{b} \cdot \vec{c})$$

$$|\vec{c} \times (\vec{a} \times \vec{b})| = |(\vec{c} \cdot \vec{b})\vec{a} - (\vec{c} \cdot \vec{a})\vec{b}|$$

$$= |(\vec{c} \cdot \vec{b})\vec{a} - \alpha(\vec{b} \cdot \vec{c})\vec{b}|$$

$$= |(\vec{c} \cdot \vec{b})| |\vec{a} - \alpha \vec{b}|$$

$$= 12 \times (|\vec{n}|)$$

$$= 12 \times 1$$

$$= 12$$

- 70.** The number of points on the curve  $y = 54x^5 - 135x^4 - 70x^3 + 180x^2 + 210x$  at which the normal lines are parallel to  $x + 90y + 2 = 0$  is :  
(1) 2                          (2) 3  
(3) 4                           (4) 0

**Official Ans. by NTA (3)**

**Ans. (3)**

**Sol.** Normal of line is parallel to line  $x + 90y + 2 = 0$

$$m_N = -\frac{1}{90}$$

$$-\left(\frac{dx}{dy}\right)_{(x_1 y_1)} = -\frac{1}{90} \Rightarrow \left(\frac{dy}{dx}\right)_{(x_1 y_1)} = 90$$

Now,

$$\frac{dy}{dx} = 270x^4 - 540x^3 - 210x^2 + 360x + 210 = 90$$

$$\Rightarrow x = 1, 2, \frac{-2}{3}, \frac{-1}{3}$$

(4) normals

- 71.** Let  $y = x + 2, 4y = 3x + 6$  and  $3y = 4x + 1$  be three tangent lines to the circle  $(x-h)^2 + (y-k)^2 = r^2$ . Then  $h+k$  is equal to :

$$(1) 5$$

$$(2) 5(1+\sqrt{2})$$

$$(3) 6$$

$$(4) 5\sqrt{2}$$

**Official Ans. by NTA (1)**

**Ans. (1)**

**Sol.**  $L_1 : y = x + 2, L_2 : 4y = 3x + 6, L_3 : 3y = 4x + 1$   
Bisector of lines  $L_2$  &  $L_3$

$$\frac{4x - 3y + 1}{5} = \pm \left( \frac{3x - 4y + 6}{5} \right)$$

$$(+) \quad 4x - 3y + 1 = 3x - 4y + 6 \\ x + y = 5$$

Centre lies on Bisector of  $4x - 3y + 1 = 0$  &

$$(0) \quad 3x - 4y + 6 = 0$$

$$\Rightarrow h + k = 5$$

- 72.** Let the solution curve  $y = y(x)$  of the differential

$$\text{equation } \frac{dy}{dx} - \frac{3x^5 \tan^{-1}(x^3)}{(1+x^6)^{\frac{3}{2}}} y = 2x$$

$\exp \frac{x^3 - \tan^{-1} x^3}{\sqrt{(1+x^6)^6}}$  pass through the origin. Then

$y(1)$  is equal to :

$$(1) \exp \left( \frac{4-\pi}{4\sqrt{2}} \right) \quad (2) \exp \left( \frac{\pi-4}{4\sqrt{2}} \right)$$

$$(3) \exp \left( \frac{1-\pi}{4\sqrt{2}} \right) \quad (4) \exp \left( \frac{4+\pi}{4\sqrt{2}} \right)$$

**Official Ans. by NTA (1)**

**Ans. (1)**

$$\text{Sol. } \frac{dy}{dx} + \left( \frac{-3x^5 \tan^{-1} x^3}{(1+x^6)^{\frac{3}{2}}} \right) y = 2e^{\frac{x - \tan^{-1} x^3}{\sqrt{1+x^6}}}$$

$$\text{I.F.} = e^{\int \frac{-3x^5 \tan^{-1} x^3}{(1+x^6)^{\frac{3}{2}}} dx} \\ = e^{\frac{\tan^{-1} x^3 - x^3}{\sqrt{1+x^6}}}$$

Solution of differential equation

$$y \cdot e^{\frac{\tan^{-1} x^3 - x^3}{\sqrt{1+x^6}}} = \int 2x e^{\left( \frac{x^3 - \tan^{-1} x^3}{\sqrt{1+x^6}} \right)} \cdot e^{\left( \frac{\tan^{-1} (x^3) - x^3}{\sqrt{1+x^6}} \right)} dx$$

$$= \int 2x dx + c$$

$$y \cdot e^{\frac{\tan^{-1} x^3 - x^3}{\sqrt{1+x^6}}} = x^2 + c$$

Also it passes through origin  
 $c = 0$

$$y(1) \cdot e^{\frac{\tan^{-1} (1)-1}{\sqrt{2}}} = 1$$

$$y(1) \cdot e^{\frac{\frac{\pi}{4}-1}{\sqrt{2}}} = 1$$

$$y(1) \cdot e^{\frac{\pi-4}{4\sqrt{2}}} = 1$$

$$y(1) = \frac{1}{e^{\frac{\pi-4}{4\sqrt{2}}}} = e^{\frac{4-\pi}{4\sqrt{2}}}$$

- 73.** Let a unit vector  $\widehat{OP}$  make angle  $\alpha, \beta, \gamma$  with the positive directions of the co-ordinate axes OX, OY, OZ respectively, where  $\beta \in \left(0, \frac{\pi}{2}\right)$   $\widehat{OP}$  is perpendicular to the plane through points (1, 2, 3), (2, 3, 4) and (1, 5, 7), then which one of the following is true ?
- $\alpha \in \left(\frac{\pi}{2}, \pi\right)$  and  $\gamma \in \left(\frac{\pi}{2}, \pi\right)$
  - $\alpha \in \left(0, \frac{\pi}{2}\right)$  and  $\gamma \in \left(0, \frac{\pi}{2}\right)$
  - $\alpha \in \left(\frac{\pi}{2}, \pi\right)$  and  $\gamma \in \left(0, \frac{\pi}{2}\right)$
  - $\alpha \in \left(0, \frac{\pi}{2}\right)$  and  $\gamma \in \left(\frac{\pi}{2}, \pi\right)$

**Official Ans. by NTA (1)**
**Ans. (1)**
**Sol.** Equation of plane :-

$$\begin{vmatrix} x-1 & y-2 & z-3 \\ 1 & 1 & 1 \\ 0 & 3 & 4 \end{vmatrix} = 0$$

$$\Rightarrow [x-1] - 4[y-2] + 3[z-3] = 0$$

$$\Rightarrow x - 4y + 3z = 2$$

D.R's of normal of plane  $\langle 1, -4, 3 \rangle$ 

D.C's of  $\left\langle \pm \frac{1}{\sqrt{26}}, \mp \frac{4}{\sqrt{26}}, \pm \frac{3}{\sqrt{26}} \right\rangle$

$$\cos \beta = \frac{4}{\sqrt{26}}$$

$$\cos \alpha = \frac{-1}{\sqrt{26}} \quad \frac{\pi}{2} < \alpha < \pi$$

$$\cos \gamma = \frac{-3}{\sqrt{26}} \quad \frac{\pi}{2} < \gamma < \pi$$

Ans. : (1)

- 74.** If [t] denotes the greatest integer  $\leq 1$ , then the value

of  $\frac{3(e-1)^2}{e} \int_1^2 x^2 e^{[x]+[x^3]} dx$  is :

- $e^9 - e$
- $e^8 - e$
- $e^7 - 1$
- $e^8 - 1$

**Official Ans. by NTA (2)**
**Ans. (2)**

**Sol.**

$$\begin{aligned} & \int_1^2 x^2 e^{[x^3]+1} dx \\ & x^3 = t \\ & 3x^2 dx = dt \\ & = \frac{e}{3} \int_1^8 e^{[t]} dt \\ & = \frac{e}{3} \left\{ \int_1^2 e^{dt} + \int_2^3 e^{2t} dt + \dots + \int_7^8 e^{7t} dt \right\} \\ & = \frac{e}{3} (e + e^2 + \dots + e^7) \\ & = \frac{e^2}{3} (1 + e + \dots + e^6) = \frac{e^2}{3} \frac{(e^7 - 1)}{(e - 1)} \\ & \frac{3(e-1)}{e} \int_1^2 x^2 \times e^{[x]+[x^3]} dx = \frac{3}{e} (e-1) \times \frac{e^2}{3} \frac{(e^7 - 1)}{(e - 1)} \\ & = e(e^7 - 1) \\ & = e^8 - e \end{aligned}$$

Ans. : (2)

- 75.** If P(h,k) be point on the parabola  $x = 4y^2$ , which is nearest to the point Q(0,33), then the distance of P from the directrix of the parabola  $y^2 = 4(x+y)$  is equal to :

- 2
- 4
- 8
- 6

**Official Ans. by NTA (4)**
**Ans. (4)**
**Sol.** Equation of normal

$$y = -tx + 2at + at^3$$

$$y = -tx + \frac{2}{16}t + \frac{1}{16}t^3$$

It passes through (0, 33)

$$33 = \frac{t}{8} + \frac{t^3}{16}$$

$$t^3 + 2t - 528 = 0$$

$$(t-8)(t^2 + 8t + 66) = 0$$

$$t = 8$$

$$P(at^2, 2at) = \left( \frac{1}{16} \times 64, 2 \times \frac{1}{16} \times 8 \right) = (4, 1)$$

Parabola :

$$y^2 = 4(x+y)$$

$$\Rightarrow y^2 - 4y = 4x$$

$$\Rightarrow (y-2)^2 = 4(x+1)$$

Equation of directix :-

$$x + 1 = -1$$

$$x = -2$$

Distance of point = 6


Ans. : (4)

$A = a$  and  $OB = b$  on the positive directions of  $x$ -axis and  $y$ -axis respectively. If the perpendicular from origin  $O$  to this line makes an angle of  $\frac{\pi}{6}$  with positive direction of  $y$ -axis and the area of  $\Delta OAB$  is  $\frac{98}{3}\sqrt{3}$ , then  $a^2 - b^2$  is equal to:

- (1)  $\frac{392}{3}$       (2) 196  
(3)  $\frac{196}{3}$       (4) 98

**Official Ans. by NTA (1)**  
**Ans. (1)**

**Sol.**



$$\text{Equation of straight line : } \frac{x}{a} + \frac{y}{b} = 1$$

$$\text{Or } x \cos \frac{\pi}{3} + y \sin \frac{\pi}{3} = p$$

$$\frac{x}{2} + \frac{y\sqrt{3}}{2} = p$$

$$\frac{x}{3p} + \frac{y}{2p} = 1$$

$$\text{Comparing both : } a = 2p, b = \frac{2p}{\sqrt{3}}$$

$$\text{Now area of } \Delta OAB = \frac{1}{2} \cdot ab = \frac{98}{3} \cdot \sqrt{3}$$

$$\frac{1}{2} \cdot 2p \cdot \frac{2p}{\sqrt{3}} = \frac{98}{3} \cdot \sqrt{3}$$

$$p^2 = 49$$

$$a^2 - b^2 = 4p^2 - \frac{4p^2}{3} = \frac{2}{3}4p^2$$

$$= \frac{8}{3} \cdot 49 = \frac{392}{3}$$

- 77.** The coefficient of  $x^{301}$  in  $(1+x)^{500} + x(1+x)^{499} + x^2(1+x)^{498} + \dots + x^{500}$  is:
- (1)  ${}^{501}C_{302}$       (2)  ${}^{500}C_{301}$   
(3)  ${}^{500}C_{300}$       (4)  ${}^{501}C_{200}$

**Official Ans. by NTA (4)**

**Ans. (4)**

$$\text{Sol. } (1+x)^{500} + x(1+x)^{499} + x^2(1+x)^{498} + \dots + x^{500}$$

$$= (1+x)^{500} \cdot \left[ \frac{1 - \left( \frac{x}{1+x} \right)^{501}}{1 - \frac{x}{1+x}} \right]$$

$$= (1+x)^{500} \frac{\left( (1+x)^{501} - x^{501} \right)}{(1+x)^{501}} \cdot (1+x)$$

$$= (1+x)^{501} - x^{501}$$

Coefficient of  $x^{301}$  in  $(1+x)^{501} - x^{501}$  is given by

$${}^{501}C_{301} = {}^{501}C_{200}$$

- 78.** Among the statements:

$$(S1) \quad ((p \vee q) \Rightarrow r) \Leftrightarrow (p \Rightarrow r)$$

$$(S2) \quad ((p \vee q) \Rightarrow r) \Leftrightarrow ((p \Rightarrow r) \vee (q \Rightarrow r))$$

(1) Only (S1) is a tautology

(2) Neither (S1) nor (S2) is a tautology

(3) Only (S2) is a tautology

(4) Both (S1) and (S2) are tautologies

**Official Ans. by NTA (2)**

**Ans. (2)**

$$\text{Sol. } S_1 \equiv ((p \vee q) \Rightarrow r) \Leftrightarrow (p \Rightarrow r)$$

$$\begin{array}{ccccccccc} p & q & r & p \vee q & (p \vee q) \Rightarrow r & p \Rightarrow r & ((p \vee q) \Rightarrow r) \Leftrightarrow (p \Rightarrow r) \\ \hline T & T & T & T & T & T & T \\ T & T & F & T & F & F & T \\ T & F & T & T & T & T & T \\ F & T & T & T & T & T & T \\ T & F & F & T & F & F & T \\ F & T & F & T & F & T & T \\ F & F & F & T & T & T & T \\ F & F & F & F & T & T & T \end{array}$$

| $S_2 \equiv (p \vee q) \Rightarrow r \Leftrightarrow ((p \Rightarrow r) \vee (q \Rightarrow r))$ |   |   |                            |                   |                   |  |    |
|--|---|---|----------------------------|-------------------|-------------------|--|----|
| p  | q | r | $(p \vee q) \Rightarrow r$ | $p \Rightarrow r$ | $q \Rightarrow r$ | $(p \Rightarrow r) \vee (q \Rightarrow r)$ | S2 |
| T  | T | T | T                          | T                 | T                 | T  | T  |
| T  | T | F | F                          | F                 | F                 | T  | T  |
| T  | F | T | T                          | T                 | T                 | T  | T  |
| F  | T | T | T                          | T                 | T                 | T  | T  |
| T  | F | F | F                          | T                 | T                 | T  | F  |
| F  | T | F | F                          | F                 | F                 | T  | F  |
| F  | F | T | T                          | T                 | T                 | T  | T  |
| F  | F | F | T                          | T                 | T                 | T  | T  |

S2 → not a tautology

79. The minimum number of elements that must be added to the relation  $R = \{(a,b), (b,c)\}$  on the set  $\{a, b, c\}$  so that it becomes symmetric and transitive is:  
(1) 4                          (2) 7  
(3) 5                          (4) 3

**Official Ans. by NTA (2)**

**Ans. (2)**

**Sol.** For Symmetric  $(a,b), (b,c) \in R$

$$\Rightarrow (b,a), (c,b) \in R$$

For Transitive  $(a,b), (b,c) \in R$

$$\Rightarrow (a,c) \in R$$

Now

1. Symmetric

$$\therefore (a,c) \in R \Rightarrow (c,a) \in R$$

2. Transitive

$$\therefore (a,b), (b,a) \in R$$

$$\Rightarrow (a,a) \in R \text{ & } (b,c), (c,b) \in R$$

$$\Rightarrow (b,b) \& (c,c) \in R$$

$\therefore$  Elements to be added

$$\left\{ (b,a), (c,b), (a,c), (c,a), (a,a), (b,b), (c,c) \right\}$$

Number of elements to be added = 7

80. If the solution of the equation  $\log_{\cos x} \cot x + 4 \log_{\sin x} \tan x = 1, x \in \left(0, \frac{\pi}{2}\right)$ , is  $\sin^{-1} \left( \frac{\alpha + \sqrt{\beta}}{2} \right)$ , where  $\alpha, \beta$  are integers, then  $\alpha + \beta$  is equal to:  
(1) 3  
(2) 5  
(3) 6  
(4) 4

**Official Ans. by NTA (4)**

**Ans. (4)**

**Sol.**

$$\begin{aligned} \log_{\cos x} \cot x + 4 \log_{\sin x} \tan x &= 1 \\ \Rightarrow \frac{\ln \cot x - \ln \cos x}{\ln \cos x} + 4 \frac{\ln \tan x - \ln \sin x}{\ln \sin x} &= 1 \\ \Rightarrow (\ln \cot x)^2 - 4(\ln \cot x)(\ln \cos x) + 4(\ln \cos x)^2 &= 1 \\ \Rightarrow \ln \cot x = 2 \ln \cos x & \\ \Rightarrow \sin^2 x + \sin x - 1 &= 0 \Rightarrow \sin x = \frac{-1 + \sqrt{5}}{2} \\ \therefore \alpha + \beta &= 4 \\ \text{Correct option (4)} & \end{aligned}$$

## SECTION-B

81. Let  $S = \{1, 2, 3, 4, 5, 6\}$ . Then the number of one-one functions  $f : S \rightarrow P(S)$ , where  $P(S)$  denote the power set of  $S$ , such that  $f(n) \subset f(m)$  where  $n < m$  is \_\_\_\_\_.

**Official Ans. by NTA (3240)**

**Ans. (3240)**

- Sol.** Let  $S = \{1, 2, 3, 4, 5, 6\}$ , then the number of one-one functions,  $f : S \rightarrow P(S)$ , where  $P(S)$  denotes the power set of  $S$ , such that  $f(n) \subset f(m)$  where  $n < m$  is

$$n(S) = 6$$

$$P(S) = \left\{ \phi, \{1\}, \dots, \{6\}, \{1, 2\}, \dots, \{5, 6\}, \dots, \{1, 2, 3, 4, 5, 6\} \right\}$$

- 64 elements

case – 1

$f(6) = S$  i.e. 1 option,

$f(5) = \text{any 5 element subset A of } S \text{ i.e. 6 options,}$

$f(4) = \text{any 4 element subset B of A i.e. 5 options,}$

$f(3) = \text{any 3 element subset C of B i.e. 4 options,}$

$f(2) = \text{any 2 element subset D of C i.e. 3 options,}$

$f(1) = \text{any 1 element subset E of D or empty subset i.e. 3 options,}$

Total functions = 1080

Case – 2

$f(6) = \text{any 5 element subset A of } S \text{ i.e. 6 options,}$

$f(5) = \text{any 4 element subset B of A i.e. 5 options,}$

$f(4) = \text{any 3 element subset C of B i.e. 4 options,}$

$f(3) = \text{any 2 element subset D of C i.e. 3 options,}$

$f(2) = \text{any 1 element subset E of D i.e. 2 options,}$

$f(1) = \text{empty subset i.e. 1 option}$

Total functions = 720

Case – 3

$f(6) = S$

$f(5) = \text{any 4 element subset A of } S \text{ i.e. 15 options,}$

$f(4) = \text{any 3 element subset B of A i.e. 4 options,}$

$f(3) = \text{any 2 element subset C of B i.e. 3 options,}$

$f(2) = \text{any 1 element subset D of C i.e. 2 options,}$

$f(1) = \text{empty subset i.e. 1 option}$

Total functions = 360

Case – 4

$f(6) = S$

$f(5) = \text{any 5 element subset A of } S \text{ i.e. 6 options,}$

$f(4) = \text{any 4 element subset B of A i.e. 10 options,}$

$f(3) = \text{any 3 element subset C of B i.e. 10 options,}$

$f(2) = \text{any 2 element subset D of C i.e. 2 options,}$

$f(1) = \text{empty subset i.e. 1 option}$

Total functions = 360

Case – 5

$f(6) = S$

$f(5) = \text{any 5 element subset A of } S \text{ i.e. 6 options,}$

$f(4) = \text{any 4 element subset B of A i.e. 5 options,}$

$f(3) = \text{any 3 element subset C of B i.e. 10 options,}$

$f(2) = \text{any 2 element subset D of C i.e. 2 options,}$

$f(1) = \text{empty subset i.e. 1 option}$

Total functions = 360

Case – 6

$f(6) = S$

$f(5) = \text{any 5 element subset A of } S \text{ i.e. 6 options,}$

$f(4) = \text{any 4 element subset B of A i.e. 5 options,}$

$f(3) = \text{any 3 element subset C of B i.e. 4 options,}$

$f(2) = \text{any 2 element subset D of C i.e. 3 options,}$

$f(1) = \text{empty subset i.e. 1 option}$

Total functions = 360


$\therefore$  Number of such functions = 3240

82. Let  $\alpha$  be the area of the larger region bounded by the curve  $y^2 = 8x$  and the lines  $y = x$  and  $x = 2$ , which lies in the first quadrant. Then the value of  $3\alpha$  is equal to \_\_\_\_\_.

Official Ans. by NTA (22)

Ans. (22)

Sol.



$$y = x$$

$$\& y^2 = 8x$$

Solving it

$$x^2 = 8x$$

$$\therefore x = 0, 8$$

$$\therefore y = 0, 8$$

$x = 2$  will intersect occur at

$$y^2 = 16 \Rightarrow y = \pm 4$$

$\therefore$  Area of shaded

$$= \int_2^8 (\sqrt{8x} - x) dx = \int_2^8 (2\sqrt{2}\sqrt{x} - x) dx$$

$$= \left[ 2\sqrt{2} \cdot \frac{x^{3/2}}{3/2} - \frac{x^2}{2} \right]_0^8$$

$$= \left( \frac{4\sqrt{2}}{3} \cdot 2^{9/2} - 32 \right) - \left( \frac{4\sqrt{2}}{3} \cdot 2^{9/2} - 2 \right)$$

$$= \frac{128}{3} - 32 - \frac{16}{3} + 2 = \frac{112 - 90}{3} = \frac{22}{3} = A$$

$$\therefore 3A = 22$$

$<\lambda$  are two values of  $\lambda$  such that the angle between the planes  $P_1 : \vec{r} \cdot (3\hat{i} - 5\hat{j} + \hat{k}) = 7$  and  $P_2 : \vec{r} \cdot (\lambda\hat{i} + \hat{j} - 3\hat{k}) = 9$  is  $\sin^{-1}\left(\frac{2\sqrt{6}}{5}\right)$ , then the square of the length of perpendicular from the point  $(38\lambda_1, 10\lambda_2, 2)$  to the plane  $P_1$  is \_\_\_\_.

**Official Ans. by NTA (315)**

**Ans. (315)**

$$\text{Sol. } P_1 = \vec{r} \cdot (3\hat{i} - 5\hat{j} + \hat{k}) = 7$$

$$P_2 = \vec{r} \cdot (\lambda\hat{i} + \hat{j} - 3\hat{k}) = 9$$

$$\theta = \sin^{-1}\left(\frac{2\sqrt{6}}{5}\right)$$

$$\Rightarrow \sin \theta = \frac{2\sqrt{6}}{5}$$

$$\therefore \cos \theta = \frac{1}{5}.$$

$$\cos \theta = \frac{\vec{r} \cdot \vec{r}}{|\vec{r}_1| |\vec{r}_2|}$$

$$= \frac{(3\hat{i} - 5\hat{j} + \hat{k})(\lambda\hat{i} + \hat{j} - 3\hat{k})}{\sqrt{35} \cdot \sqrt{\lambda^2 + 10}}$$


$$\frac{1}{5} = \left| \frac{3\lambda - 8}{\sqrt{35} \cdot \sqrt{\lambda^2 + 10}} \right|$$

$$\text{Square} \Rightarrow \frac{1}{25} = \frac{9\lambda^2 + 64 - 48\lambda}{35(\lambda^2 + 10)}$$

$$\Rightarrow 19\lambda^2 - 120\lambda + 125 = 0$$

$$\Rightarrow 19\lambda^2 - 95\lambda - 25\lambda + 125 = 0$$

$$\Rightarrow x = 5, \frac{25}{19}$$



Perpendicular distance of point

$$(38\lambda_1, 10\lambda_2, 2) \equiv (50, 50, 2) \text{ from plane } P_1$$

$$= \frac{|3 \times 50 - 5 \times 50 + 2 - 7|}{\sqrt{35}} = \frac{105}{\sqrt{35}}$$

$$\text{Square} = \frac{105 \times 105}{35} = 315$$

84. Let  $z = 1+i$  and  $z_1 = \frac{1+i\bar{z}}{\bar{z}(1-z)+\frac{1}{z}}$ . Then  $\frac{12}{\pi} \arg(z_1)$  is equal to \_\_\_\_.

**Official Ans. by NTA (9)**

**Ans. (9)**

$$\text{Sol. } z = 1+i$$

$$z_1 = \frac{1+i\bar{z}}{\bar{z}(1-z)+\frac{1}{z}}$$

$$z_1 = \frac{1+i(1-i)}{(1-i)(1-1-i)+\frac{1}{1+i}}$$

$$= \frac{1+i-i^2}{(1-i)(-i)+\frac{1-i}{2}}$$

$$= \frac{2+i}{-3i-1} = \frac{4+2i}{-3i-1}$$

$$= \frac{-(4+2i)(3i-1)}{(3i)^2 - (1)^2}$$

$$\text{Arg}(z_1) = \frac{3\pi}{4}$$

$$\therefore \frac{12}{\pi} \arg(z_1) = \frac{12}{\pi} \times \frac{3\pi}{4} = 9$$

85.  $\lim_{x \rightarrow 0} \frac{48 \int_0^x \frac{t^3}{t^6 + 1} dt}{x^4}$  is equal to \_\_\_\_\_.

**Official Ans. by NTA (12)**

**Ans. (12)**

Sol.  $48 \lim_{x \rightarrow 0} \frac{\int_0^x \frac{t^3}{t^6 + 1} dt}{x^4} \left( \frac{0}{0} \right)$

Applying L' Hospitals Rule

$$48 \lim_{x \rightarrow 0} \frac{x^3}{x^6 + 1} \times \frac{1}{4x^3}$$

$$= 12$$

86. The mean and variance of 7 observations are 8 and 16 respectively. If one observation 14 is omitted a and b are respectively mean and variance of remaining 6 observation, then  $a + 3b - 5$  is equal to \_\_\_\_\_.

**Official Ans. by NTA (37)**

**Ans. (37)**

Sol.  $\frac{x_1 + x_2 + \dots + x_7}{7} = 8$

$$\frac{x_1 + x_2 + x_3 + \dots + x_6 + 14}{7} = 8$$

$$\Rightarrow x_1 + x_2 + \dots + x_6 = 42$$

$$\therefore \frac{x_1 + x_2 + \dots + x_6}{6} = \frac{42}{6} = 7 = a$$

$$\frac{\sum x_i^2}{7} - 8^2 = 16$$

$$\Sigma x_i^2 = 560$$

$$\Rightarrow x_1^2 + x_2^2 + \dots + x_6^2 = 364$$

$$b = \frac{x_1^2 + x_2^2 + \dots + x_6^2}{6} - 7^2$$

$$= \frac{364}{6} - 49$$

$$b = \frac{70}{6}$$

$$a + 3b - 5 = 7 + 3 \times \frac{70}{6} - 5$$

$$= 37$$

87. If the equation of the plane passing through the point  $(1, 1, 2)$  and perpendicular to the line  $x - 3y + 2z - 1 = 0$   $4x - y + z$  is  $Ax + By + Cz = 1$ , then  $140(C - B + A)$  is equal to \_\_\_\_\_.

**Official Ans. by NTA (15)**

**Ans. (15)**

Sol.  $x - 3y + 2z - 1 = 0$

$$4x - y + z = 0$$

$$\vec{n}_1 \times \vec{n}_2 = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & -3 & 2 \\ 4 & -1 & 1 \end{vmatrix}$$

$$= -\hat{i} + 7\hat{j} + 11\hat{k}$$

Dr<sup>s</sup> of normal to the plane is  $-1, 7, 11$

Equation of plane :

$$-1(x - 1) + 7(y - 1) + 11(z - 2) = 0$$

$$-x + 7y + 11z = 28$$

$$\frac{-1}{28}x + \frac{7y}{28} + \frac{11z}{28} = 1$$

$$Ax + By + Cz = 1$$

$$140(C - B + A) = 140 \left( \frac{11}{28} - \frac{7}{28} - \frac{1}{28} \right)$$

$$= 140 \times \frac{3}{28} = 15$$

88. Let  $\sum_{n=0}^{\infty} \frac{n^3((2n)!) + (2n-1)(n)!}{(n!)((2n)!)}$   $= ae + \frac{b}{e} + c$ ,

where  $a, b, c \in \mathbb{Z}$  and  $e = \sum_{n=0}^{\infty} \frac{1}{n!}$ . Then  $a^2 - b + c$  is

equal to \_\_\_\_\_.

**Official Ans. by NTA (26)**

**Ans. (26)**

**Sol.** 
$$\sum_{n=0}^{\infty} \frac{n^3((2n)!) + (2n-1)(n!)}{(n!)((2n)!)}$$

$$= \sum_{n=0}^{\infty} \frac{1}{(n-3)!} + \sum_{n=0}^{\infty} \frac{3}{(n-2)!}$$

$$+ \sum_{n=0}^{\infty} \frac{1}{(n-1)!} + \sum_{n=0}^{\infty} \frac{1}{(2n-1)!} - \sum_{n=0}^{\infty} \frac{1}{(2n)!}$$

$$= e + 3e + e + \frac{1}{2} \left( e - \frac{1}{e} \right) - \frac{1}{2} \left( e + \frac{1}{e} \right)$$

$$= 5e - \frac{1}{e}$$

$$\therefore a^2 - b + c = 26$$

- 89.** Number of 4-digit numbers (the repetition of digits is allowed) which are made using the digits 1, 2, 3 and 5, and are divisible by 15, is equal to \_\_\_\_\_

**Official Ans. by NTA (21)**

**Ans. (21)**

**Sol.** For number to be divisible by 15, last digit should be 5 and sum of digits must be divisible by 3.

Possible combinations are

|   |   |   |   |
|---|---|---|---|
| 1 | 2 | 1 | 5 |
|---|---|---|---|

Numbers = 3

|   |   |   |   |
|---|---|---|---|
| 2 | 2 | 3 | 5 |
|---|---|---|---|

Numbers = 3

|   |   |   |   |
|---|---|---|---|
| 3 | 3 | 1 | 5 |
|---|---|---|---|

Numbers = 3

|   |   |   |   |
|---|---|---|---|
| 1 | 1 | 5 | 5 |
|---|---|---|---|

Numbers = 3

|   |   |   |   |
|---|---|---|---|
| 2 | 3 | 5 | 5 |
|---|---|---|---|

Numbers = 6

|   |   |   |   |
|---|---|---|---|
| 3 | 5 | 5 | 5 |
|---|---|---|---|

Numbers = 3

Total Numbers = 21

**90.** Let  $f^1(x) = \frac{3x+2}{2x+3}, x \in \mathbb{R} - \left\{ \frac{-3}{2} \right\}$

For  $n \geq 2$ , define  $f^n(x) = f^{10f^{n-1}(x)}$ .

If  $f^5(x) = \frac{ax+b}{bx+a}$ ,  $\gcd(a,b)=1$ , then  $a+b$  is equal to \_\_\_\_\_.

**Official Ans. by NTA (3125)**

**Ans. (3125)**

**Sol.**  $f^1(x) = \frac{3x+2}{2x+3}$

$$\Rightarrow f^2(x) = \frac{13x+12}{12x+13}$$

$$\Rightarrow f^3(x) = \frac{63x+62}{62x+63}$$

$$\therefore f^5(x) = \frac{1563x+1562}{1562x+1563}$$

$$a+b = 3125$$