



FINAL JEE–MAIN EXAMINATION – APRIL, 2019

Held On Wednesday 10th APRIL, 2019

TIME: 2 : 30 PM To 5 : 30 PM

1. The correct match between Item-I and Item-II is:

| | Item-I | | Item-II |
|-----|------------------------|-------|---|
| (a) | High density polythene | (I) | Peroxide catalyst |
| (b) | Polyacrylonitrile | (II) | Condensation at high temperature & pressure |
| (c) | Novolac | (III) | Ziegler-Natta Catalyst |
| (d) | Nylon 6 | (IV) | Acid or base catalyst |

- (1) (a)→(III), (b)→(I), (c)→(II), (d)→(IV)
 (2) (a)→(IV), (b)→(II), (c)→(I), (d)→(III)
 (3) (a)→(II), (b)→(IV), (c)→(I), (d)→(III)
 (4) (a)→(III), (b)→(I), (c)→(IV), (d)→(II)

Official Ans. by NTA (4)

Sol.



| | | | |
|-----|------------------------|-------|---|
| (a) | High density polythene | (III) | Ziegler-Natta Catalyst |
| (b) | Polyacrylonitrile | (I) | Peroxide catalyst |
| (c) | Novolac | (IV) | Acid or base catalyst |
| (d) | Nylon 6 | (II) | Condensation at high temperature & pressure |

2. Which of the following is NOT a correct method of the preparation of benzylamine from cyanobenzene ?

- (1) (i) HCl/H₂O (ii) NaBH₄
 (2) (i) LiAlH₄ (ii) H₃O⁺
 (3) (i) SnCl₂+HCl(gas) (ii) NaBH₄
 (4) H₂/Ni

Official Ans. by NTA (1)

Sol.



3. Which of these factors does not govern the stability of a conformation in acyclic compounds ?

- (1) Torsional strain
 (2) Angle strain
 (3) Steric interactions
 (4) Electrostatic forces of interaction

Official Ans. by NTA (2)

Sol. in acyclic compounds angle strain does not govern the stability of a conformation.

4. The difference between ΔH and ΔU ($\Delta H - \Delta U$), when the combustion of one mole of heptane (1) is carried out at a temperature T, is equal to:

- (1) 3RT (2) -3RT (3) -4RT (4) 4RT

Official Ans. by NTA (3)

Sol. $C_7H_{16}(l) + 11O_2(g) \longrightarrow 7CO_2(g) + 8H_2O(l)$
 $\Delta n_g = n_p - n_r = 7 - 11 = -4$

$$\therefore \Delta H = \Delta U + \Delta n_g RT$$

$$\therefore \Delta H - \Delta U = -4 RT$$

5. For the reaction of H₂ with I₂, the rate constant is $2.5 \times 10^{-4} \text{ dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$ at 327°C and $1.0 \text{ dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$ at 527°C. The activation energy for the reaction, in kJ mol⁻¹ is:

$$(R = 8.314 \text{ J K}^{-1} \text{ mol}^{-1})$$

- (1) 72 (2) 166 (3) 150 (4) 59

Official Ans. by NTA (2)

Sol. $H_2(g) + I_2(g) \rightarrow 2HI(g)$

Apply Arrhenius equation




$$\log \frac{K_2}{K_1} = \frac{E_a}{2.303R} \left(\frac{1}{600} - \frac{1}{800} \right)$$

$$\log \frac{1}{2.5 \times 10^{-4}} = \frac{E_a}{2.303 \times 8.31} \left(\frac{200}{600 \times 800} \right)$$

$$\therefore E_a \approx 166 \text{ kJ/mol}$$

6. The correct statements among (a) to (b) are:
- saline hydrides produce H_2 gas when reacted with H_2O .
 - reaction of $LiAlH_4$ with BF_3 leads to B_2H_6 .
 - PH_3 and CH_4 are electron - rich and electron-precise hydrides, respectively.
 - HF and CH_4 are called as molecular hydrides.
- (1) (c) and (d) only
 (2) (a), (b) and (c) only
 (3) (a), (b), (c) and (d)
 (4) (a), (c) and (d) only

Official Ans. by NTA (3)




- (d) HF & CH_4 are molecular hydride due to they are covalent molecules.

7. The increasing order of nucleophilicity of the following nucleophiles is :



- (1) (b) < (c) < (a) < (d)
 (2) (a) < (d) < (c) < (b)
 (3) (d) < (a) < (c) < (b)
 (4) (b) < (c) < (d) < (a)


Official Ans. by NTA (1)




ione pair donating tendency on oxygen is reduced, nucleophilicity reduced $b < c < a < d$

8. Number of stereo centers present in linear and cyclic structures of glucose are respectively :
- (1) 4 & 5 (2) 5 & 5 (3) 4 & 4 (4) 5 & 4
- Official Ans. by NTA (1)**

Sol.



D-Glucose
(Linear structure)




α-D-Glucose
(cyclic structure)

* :- Stereocenter

9. A hydrated solid X on heating initially gives a monohydrated compound Y. Y upon heating above 373K leads to an anhydrous white powder Z. X and Z, respectively, are:
- Washing soda and soda ash.
 - Washing soda and dead burnt plaster.
 - Baking soda and dead burnt plaster.
 - Baking soda and soda ash.

Official Ans. by NTA (1)



10. The number of pentagons in C_{60} and trigons (triangles) in white phosphorus, respectively, are:
- 12 and 3
 - 20 and 4
 - 12 and 4
 - 20 and 3

Official Ans. by NTA (3)

Sol. Total No. of pentagons in $C_{60} = 12$
 Total no. of trigons (triangles) in white phosphorus (P_4) = 4

11. The correct order of the first ionization enthalpies is:
- $Mn < Ti < Zn < Ni$
 - $Ti < Mn < Ni < Zn$
 - $Zn < Ni < Mn < Ti$
 - $Ti < Mn < Zn < Ni$

Official Ans. by NTA (2)




Sol. Ti \rightarrow [Ar] 3d² 4s²
 Mn \rightarrow [Ar] 3d⁵ 4s²
 Ni \rightarrow [Ar] 3d⁸ 4s²
 Zn \rightarrow [Ar] 3d¹⁰ 4s²
 Correct order of I.P. is
 [Ti < Mn < Ni < Zn]

- 12.** The correct option among the following is :
- (1) Colloidal particles in lyophobic sols can be precipitated by electrophoresis.
 - (2) Brownian motion in colloidal solution is faster the viscosity of the solution is very high.
 - (3) Colloidal medicines are more effective because they have small surface area.
 - (4) Addition of alum to water makes it unfit for drinking.

Official Ans. by NTA (1)

Sol. In electrophoresis precipitation occurs at the electrode which is oppositely charged therefore (1) is correct.

- 13.** Points I, II and III in the following plot respectively correspond to
 (V_{mp} : most probable velocity)



- (1) V_{mp} of N₂ (300K); V_{mp} of H₂(300K); V_{mp} of O₂(400K)
- (2) V_{mp} of H₂ (300K); V_{mp} of N₂(300K); V_{mp} of O₂(400K)
- (3) V_{mp} of O₂ (400K); V_{mp} of N₂(300K); V_{mp} of H₂(300K)
- (4) V_{mp} of N₂ (300K); V_{mp} of O₂(400K); V_{mp} of H₂(300K)

Official Ans. by NTA (4)

Sol. $V_{mp} = \sqrt{\frac{2RT}{M}} \Rightarrow V_{mp} \propto \sqrt{\frac{T}{M}}$

For N₂, O₂, H₂

$$\sqrt{\frac{300}{28}} < \sqrt{\frac{400}{32}} < \sqrt{\frac{300}{2}}$$

$$V_{mp} \text{ of N}_2(300K) < V_{mp} \text{ of O}_2(400K) < V_{mp} \text{ of H}_2(300K)$$

14. The INCORRECT statement is :

- (1) the spin-only magnetic moments of [Fe(H₂O)₆]²⁺ and [Cr(H₂O)₆]²⁺ are nearly similar.
- (2) the spin-only magnetic moment of [Ni(NH₃)₄(H₂O)₂]²⁺ is 2.83BM.
- (3) the gemstone, ruby, has Cr³⁺ ions occupying the octahedral sites of beryl.
- (4) the color of [CoCl(NH₃)₅]²⁺ is violet as it absorbs the yellow light.

Official Ans. by NTA (3)

- Sol.** (1) [Fe(H₂O)₆]²⁺, Fe²⁺ \rightarrow 3d⁶ \rightarrow 4 unpaired electron
 [Cr(H₂O)₆]²⁺, Cr²⁺ \rightarrow 3d⁴ \rightarrow 4 unpaired electron
 (2) [Ni(NH₃)₄(H₂O)₂]²⁺ = Ni²⁺ \rightarrow 3d⁸
 \rightarrow 2 unpaired electron
 $\mu_m = 2.83 \text{ B.M}$
 (3) In gemstone, ruby has Cr³⁺ ion occupying the octahedral sites of aluminium oxide (Al₂O₃) normally occupied by Al³⁺ ion.
 (4) Complimentary color of violet is yellow

- 15.** For the reaction,
 2SO₂(g) + O₂(g) \rightleftharpoons 2SO₃(g),
 $\Delta H = -57.2 \text{ kJ mol}^{-1}$ and
 $K_c = 1.7 \times 10^{16}$.
 Which of the following statement is INCORRECT?

- (1) The equilibrium constant is large suggestive of reaction going to completion and so no catalyst is required.
- (2) The equilibrium will shift in forward direction as the pressure increase.
- (3) The equilibrium constant decreases as the temperature increases.
- (4) The addition of inert gas at constant volume will not affect the equilibrium constant.

Official Ans. by NTA (1)

- Sol.** In option (2)- Δn_g is -ve therefore increase in pressure will bring reaction in forward direction.
 In option (3)- as the reaction is exothermic therefore increase in temperature will decrease the equilibrium constant.
 In option (4)- Equilibrium constant changes only with temperature.
 Hence, option (2), (3) and (4) are correct therefore option (1) is incorrect choice.



16. The pH of a 0.02M NH_4Cl solution will be
 [given $K_b(\text{NH}_4\text{OH})=10^{-5}$ and $\log 2=0.301$]
 (1) 4.65 (2) 5.35
 (3) 4.35 (4) 2.65

Official Ans. by NTA (2)

Sol. For the salt of strong acid and weak base

$$[\text{H}^+] = \sqrt{\frac{K_w \times C}{K_b}}$$

$$[\text{H}^+] = \sqrt{\frac{10^{-14} \times 2 \times 10^{-2}}{10^{-5}}}$$

$$-\log[\text{H}^+] = 6 - \frac{1}{2} \log 20$$

$$\therefore \text{pH} = 5.35$$

17. The noble gas that does NOT occur in the atmosphere is:
 (1) He (2) Ra
 (3) Ne (4) Kr

Official Ans. by NTA (2)

Ans. (Bonus)

Sol. In question noble gas asked, which does not exist in the atmosphere and answer is given Ra. Ra is a alkaline earth metal not noble gas it should be Rn. It is printing error in JEE Main paper

18. 1 g of non-volatile non-electrolyte solute is dissolved in 100g of two different solvents A and B whose ebullioscopic constants are in the ratio of 1 : 5. The ratio of the elevation in their boiling

points, $\frac{\Delta T_b(\text{A})}{\Delta T_b(\text{B})}$, is :

- (1) 5 : 1
 (2) 10 : 1
 (3) 1 : 5
 (4) 1 : 0.2





Official Ans. by NTA (3)

Sol. $\Delta T_b = K_b \times m$

$$\therefore \frac{\Delta T_b(\text{A})}{\Delta T_b(\text{B})} = \frac{K_b(\text{A})}{K_b(\text{B})} \text{ as } m_A = m_B$$

$$\therefore \frac{\Delta T_b(\text{A})}{\Delta T_b(\text{B})} = \frac{1}{5}$$

19. Which one of the following graphs between molar conductivity (Λ_m) versus \sqrt{C} is correct?



Official Ans. by NTA (2)

Sol. Both NaCl and KCl are strong electrolytes and as $\text{Na}^+(\text{aq.})$ has less conductance than $\text{K}^+(\text{aq.})$ due to more hydration therefore the graph of option (2) is correct.



20. The correct statement is :
- (1) zincite is a carbonate ore
 - (2) aniline is a froth stabilizer
 - (3) zone refining process is used for the refining of titanium
 - (4) sodium cyanide cannot be used in the metallurgy of silver

Official Ans. by NTA (2)

- Sol.** (1) Zincite is ZnO
 (2) Aniline is the forth stablizer.
 (3) Zone refining process is not used for refining of 'Ti'
 (4) Sodium cyanide is used in the metallurgy of silver

21. The minimum amount of O₂(g) consumed per gram of reactant is for the reaction :

(Given atomic mass : Fe = 56, O = 16, Mg = 24, P = 31, C = 12, H = 1)

- (1) $C_3H_8(g) + 5 O_2(g) \rightarrow 3 CO_2(g) + 4 H_2O(l)$
- (2) $P_4(s) + 5 O_2(g) \rightarrow P_4O_{10}(s)$
- (3) $4 Fe(s) + 3 O_2(g) \rightarrow 2 Fe_2O_3(s)$
- (4) $2 Mg(s) + O_2(g) \rightarrow 2 MgO(s)$

Official Ans. by NTA (3)

- Sol.** $C_3H_8(g) + 5O_2(g) \longrightarrow 3CO_2(g) + 4H_2O(l)$

Each 1g of C₃H₈ requires 3.63 g of O₂

$P_4(s) + 5O_2(g) \longrightarrow P_4O_{10}(s)$

Each 1g of P₄ requires 1.29 g of O₂

$4Fe(s) + 3O_2(g) \longrightarrow 2Fe_2O_3(s)$

Each 1g of Fe requires 0.428 g of O₂

$2Mg(s) + O_2(g) \longrightarrow 2MgO(s)$

Each 1g of Mg requires 0.66 g of O₂

therefore least amount of O₂ is required in option (3).


22. Air pollution that occurs in sunlight is :

- (1) oxidising smog
- (2) acid rain
- (3) reducing smog
- (4) fog

Official Ans. by NTA (1)


- Sol.** Photochemical smog occurs in warm (sunlight) and has high concentration of oxidising agent therefore it is called photochemical smog/oxidising smog.

23. The major product 'Y' in the following reaction is:



- (1)
- (2)
- (3)
- (4)

Official Ans. by NTA (3)




24. Compound A (C₉H₁₀O) shows positive iodoform test. Oxidation of A with KMnO₄/KOH gives acid B (C₈H₆O₄). Anhydride of B is used for the preparation of phenolphthalein. Compound A is :-


- (1)
- (2)
- (3)
- (4)

Official Ans. by NTA (1)

Sol.



+ve iodoform test



Pthleic anhydride

is used for preparation of phenolphthalein indicator




25. The crystal field stabilization energy (CFSE) of $[\text{Fe}(\text{H}_2\text{O})_6]\text{Cl}_2$ and $\text{K}_2[\text{NiCl}_4]$, respectively, are :-

- (1) $-0.4\Delta_o$ and $-0.8\Delta_t$
- (2) $-0.4\Delta_o$ and $-1.2\Delta_t$
- (3) $-2.4\Delta_o$ and $-1.2\Delta_t$
- (4) $-0.6\Delta_o$ and $-0.8\Delta_t$

Official Ans. by NTA (1)


Sol. $[\text{Fe}(\text{H}_2\text{O})_6]\text{Cl}_2$, $\text{Fe}^{2+} \rightarrow 3d^6 \rightarrow (t_2g)^4(e_g)^2$
 C.F.S.E. = $4 \times (-0.4\Delta_o) + 2 \times 0.6\Delta_o = -0.4\Delta_o$
 $\text{K}_2[\text{NiCl}_4]$, $\text{Ni}^{2+} \rightarrow 3d^8 \rightarrow (e)^4(t_2)^4$
 C.F.S.E. = $4 \times (-0.6\Delta_t) + 4 \times (0.4\Delta_t) = -0.8\Delta_t$

26. The major product obtained in the given reaction is :-



- (1)
- (2)
- (3)
- (4)

Official Ans. by NTA (4)



27. The highest possible oxidation states of uranium and plutonium, respectively, are :-

- (1) 6 and 4
- (2) 7 and 6
- (3) 4 and 6
- (4) 6 and 7

Official Ans. by NTA (4)

Sol. The highest oxidation state of U and Pu is 6+ and 7+ respectively


28. In chromatography, which of the following statements is INCORRECT for R_f ?

- (1) R_f value depends on the type of chromatography.
- (2) The value of R_f can not be more than one.
- (3) Higher R_f value means higher adsorption.
- (4) R_f value is dependent on the mobile phase.

Official Ans. by NTA (3)


Sol. Except (3) all are correct

29. The major product 'Y' in the following reaction is:-



- (1)
- (2)
- (3)
- (4)

Official Ans. by NTA (1)



30. The ratio of the shortest wavelength of two spectral series of hydrogen spectrum is found to be about 9. The spectral series are:

- (1) Paschen and P fund
- (2) Lyman and Paschen
- (3) Brackett and Piund
- (4) Balmer and Brackett

Official Ans. by NTA (2)

Sol.

$$\frac{1}{\lambda_2} = R_H \left(\frac{1}{n_1^2} - \frac{1}{n_2^2} \right) Z^2$$

$$\frac{1}{\lambda_1} = R_H \left(\frac{1}{m_1^2} - \frac{1}{m_2^2} \right) Z^2$$

as for shortest wavelengths both n_2 and m_2 are ∞

$$\therefore \frac{\lambda_1}{\lambda_2} = \frac{9}{1} = \frac{m_1^2}{n_1^2}$$

Now if $m_1 = 3$ & $n_1 = 1$ it will justify the statement hence Lyman and Paschen (2) is correct.